

Sun Pharmaceutical Industries Limited

Sun House, Plot No. 201 B/1,
Western Express Highway, Goregaon (E),
Mumbai – 400 063, Maharashtra, INDIA.
Tel.: (91-22) 4324 4324
Fax: (91-22) 4324 4343
Website: www.sunpharma.com
Email: secretarial@sunpharma.com
CIN: L24230GJ1993PLC019050



21 July 2025

National Stock Exchange of India Limited
Scrip Symbol: SUNPHARMA

BSE Limited
Scrip Code: 524715

Intimation under Regulation 30 of SEBI (Listing Obligations and Disclosure Requirements) Regulations, 2015 – Press Release

Enclosed herewith is a copy of the press release by Sun Pharmaceutical Industries Limited (together with its subsidiaries / affiliated companies, “Sun Pharma”) about “Phase 3 Clinical Studies Evaluating Tildrakizumab 100 mg (ILUMYA) in Active Psoriatic Arthritis Meet their Primary Endpoint.”

For **Sun Pharmaceutical Industries Limited**

(Anoop Deshpande)
Company Secretary and Compliance Officer
ICSI Membership No.: A23983



FOR IMMEDIATE RELEASE

Sun Pharma's Phase 3 Clinical Studies Evaluating Tildrakizumab 100 mg (ILUMYA®) in Active Psoriatic Arthritis Meet their Primary Endpoint

- *The INSPIRE-1 and INSPIRE-2 trials are Phase 3 studies assessing the efficacy and safety of tildrakizumab 100 mg (ILUMYA) in adults with active psoriatic arthritis (PsA)*
- *Both pivotal studies met their primary endpoint with statistically significant improvements based on ACR20 response rates at Week 24 with tildrakizumab 100 mg (ILUMYA) compared to placebo*
- *These results support the potential regulatory submission of ILUMYA for the treatment of active psoriatic arthritis in the US*

MUMBAI, India & PRINCETON, N.J., July 21, 2025 – Sun Pharmaceutical Industries Limited (Reuters: SUN.BO, Bloomberg: SUNP IN, NSE: SUNPHARMA, BSE: 524715 (together with its subsidiaries and/or affiliated companies, “Sun Pharma”) today announced top-line results from two Phase 3 clinical studies evaluating the efficacy and safety of tildrakizumab 100 mg (ILUMYA®) administered over 24 weeks for treatment of active psoriatic arthritis.

Treatment with tildrakizumab 100 mg (ILUMYA) resulted in greater improvements in PsA signs and symptoms at Week 24 compared to treatment with placebo. Both the INSPIRE-1 and INSPIRE-2 studies achieved the primary endpoint, with a higher proportion of patients in the INSPIRE-1 and INSPIRE-2 studies treated with tildrakizumab achieving ACR20 responses at week 24, compared to those receiving placebo ($p < 0.05$).

"We are excited to share that both the INSPIRE-1 and INSPIRE-2 clinical trials have successfully met their primary endpoints. These top-line results reinforce the therapeutic potential of ILUMYA as a treatment option for patients with active psoriatic arthritis. We extend our sincere gratitude to the patients, healthcare professionals and administrators whose contributions made the studies possible. We look forward to sharing the complete clinical data in the near future," said Marek Honczarenko, MD, PhD, Senior Vice President and Head of Global Specialty Development at Sun Pharma.

Safety data in the studies was consistent with the well-documented safety profile of ILUMYA, which is approved for the treatment of adults with moderate-to-severe plaque psoriasis who are candidates for systemic therapy or phototherapy. No new safety signals were identified in the INSPIRE-1 and INSPIRE-2 studies.

Findings from the INSPIRE studies will be presented at upcoming medical conferences and published in a peer-reviewed medical journal. Use of tildrakizumab 100 mg (ILUMYA) in psoriatic arthritis is not approved, and its safety and efficacy have not been evaluated by regulatory authorities.



About the INSPIRE-1 and INSPIRE-2 Phase 3 studies

The INSPIRE-1 and INSPIRE-2 are both 52-week global, multicenter, randomized, double-blind, single-dose, placebo-controlled Phase 3 studies aimed at assessing the efficacy and safety of tildrakizumab 100 mg (ILUMYA®) in adult patients with active psoriatic arthritis. The INSPIRE-1 enrolled patients having prior exposure to an anti-TNF agent while the INSPIRE-2 enrolled anti-TNF naïve patients. A total of over 800 adult patients were enrolled for the two studies from clinical sites in the US, Europe and Asia. Patients were randomized to receive either tildrakizumab 100 mg (ILUMYA®) or placebo. An induction dose was not administered in either study. Study participants were permitted to use concomitant methotrexate or leflunomide, provided the dose remained stable throughout the trial. In the INSPIRE-1 and INSPIRE-2 studies, tildrakizumab 100 mg (ILUMYA®) was administered at Week 0 and then every 12 weeks. In contrast, in Phase 3 studies for chronic plaque psoriasis, the treatment schedule included doses in Week 0, Week 4, and then every 12 weeks thereafter. ILUMYA® is approved by the regulatory Agencies for the indication of chronic plaques psoriasis.

The primary endpoint for both studies is the proportion of participants achieving an ACR20 response at Week 24. The key secondary efficacy endpoints at 24 weeks include ACR50, ACR70, and PASI75 and improvement. Learn more about the studies at clinicaltrials.gov (INSPIRE-1 NCT04314544 and INSPIRE-2 NCT04314531).

The ACR20 is a composite measure defined as a 20 percent improvement in both the number of tender and number of swollen joints and a 20 percent improvement in three of the following five criteria: patient global assessment, physician global assessment, functional ability measure (most often HAQ-DI), visual analog pain scale, and erythrocyte sedimentation rate or C-reactive protein (CRP). ACR50 and ACR70 are the same measurement with improvement levels of 50 percent and 70 percent, respectively.

About Psoriatic Arthritis (PsA)

Psoriatic arthritis (PsA) is a chronic (long-lasting or recurring) disease related to the immune system. It causes swelling, pain, and stiffness in joints and entheses (places where tendons and ligaments connect to bones).[1] Psoriatic arthritis can develop at any age.[2] Psoriatic arthritis can occur regardless of the severity of psoriasis (mild, moderate, or severe). Roughly 1 in 3 people living with psoriasis also have psoriatic arthritis [3]; typically, with psoriasis developing before psoriatic arthritis.[2] While as many as 2.4 million Americans live with psoriatic arthritis, more than 15% of people living with psoriasis may also have undiagnosed psoriatic arthritis.[4]

About ILUMYA® (tildrakizumab-asmn)

ILUMYA (tildrakizumab-asmn) is a humanized IgG1/k monoclonal antibody designed to selectively bind to the p19 subunit of interleukin-23 (IL-23) and inhibit its interaction with the IL-23 receptor, leading to inhibition of the release of pro-inflammatory cytokines and chemokines. ILUMYA is indicated for the treatment of adults with moderate-to-



severe plaque psoriasis who are candidates for systemic therapy or phototherapy, in the United States and other countries.

INDICATIONS AND USAGE

ILUMYA (tildrakizumab-asmn) is indicated for the treatment of adults with moderate-to-severe plaque psoriasis who are candidates for systemic therapy or phototherapy.

CONTRAINDICATIONS

ILUMYA is contraindicated in patients with a previous serious hypersensitivity reaction to tildrakizumab or to any of the excipients.

IMPORTANT SAFETY INFORMATION

WARNINGS AND PRECAUTIONS

Hypersensitivity

Cases of angioedema and urticaria occurred in ILUMYA-treated patients in clinical trials. If a serious allergic reaction occurs, discontinue ILUMYA immediately and initiate appropriate therapy.

Infections

ILUMYA may increase the risk of infection. Treatment with ILUMYA should not be initiated in patients with a clinically important active infection until the infection resolves or is adequately treated.

Consider the risks and benefits of treatment prior to prescribing ILUMYA in patients with a chronic infection or a history of recurrent infection. Instruct patients receiving ILUMYA to seek medical help if signs or symptoms of clinically important chronic or acute infection occur. If a patient develops a clinically important or serious infection, or is not responding to standard therapy, closely monitor and consider discontinuation of ILUMYA until the infection resolves.

Pretreatment Evaluation for Tuberculosis

Evaluate patients for tuberculosis (TB) infection prior to initiating treatment with ILUMYA. Do not administer ILUMYA to patients with active TB infection. Initiate treatment of latent TB prior to administering ILUMYA. Consider anti-TB therapy prior to initiation of ILUMYA in patients with a past history of latent or active TB in whom an adequate course of treatment cannot be confirmed. Patients receiving ILUMYA should be monitored closely for signs and symptoms of active TB during and after treatment.

Immunizations

Prior to initiating therapy with ILUMYA, consider completion of all age-appropriate immunizations according to current immunization guidelines. Patients treated with ILUMYA should not receive live vaccines.

Sun Pharmaceutical Industries Limited
SUN HOUSE, Plot No. 201 B/1,
Western Express Highway, Goregaon (E),
Mumbai 400063, India
Tel.: (91-22) 4324 4324 Fax.: (91-22) 4324 4343
Website: www.sunpharma.com
Email: secretarial@sunpharma.com
CIN: L24230GJ1993PLC019050



Adverse Reactions

The most common ($\geq 1\%$ adverse reactions associated with ILUMYA treatment that were more frequent than in the placebo group are upper respiratory infections, injection-site reactions, and diarrhea.

Please see [Full Prescribing Information](#) and [Medication Guide](#)

About Sun Pharmaceutical Industries Limited. (CIN - L24230GJ1993PLC019050)

Sun Pharma is the world's leading specialty generics company with a presence in specialty, generics and consumer healthcare products. It is the largest pharmaceutical company in India and is a leading generic company in the U.S. as well as global emerging markets. Sun Pharma's high-growth global specialty portfolio spans innovative products in dermatology, ophthalmology, and onco-dermatology and accounts for about 20% of company sales. The company's vertically integrated operations deliver high-quality medicines, trusted by physicians and consumers in over 100 countries. Its manufacturing facilities are spread across six continents. Sun Pharma is proud of its multicultural workforce drawn from over 50 nations. For further information, please visit www.sunpharma.com and follow us on [LinkedIn](#) & [X](#) (Formerly Twitter).

Disclaimer

Statements in this "Document" describing the Company's objectives, projections, estimates, expectations, plans or predictions or industry conditions or events may be "forward looking statements" within the meaning of applicable securities laws and regulations. Actual results, performance or achievements could differ materially from those expressed or implied. The Company undertakes no obligation to update or revise forward looking statements to reflect developments or circumstances that arise or to reflect the occurrence of unanticipated developments/circumstances after the date hereof.

References

1. National Psoriasis Foundation. About Psoriatic Arthritis. 2025.
2. Ritchlin CT, Colbert RA, Gladman DD. Psoriatic Arthritis [published correction appears in N Engl J Med. 2017 May 25;376(21):2097]. N Engl J Med. 2017;376(10):957-970. doi:10.1056/NEJMra1505557
3. Mease PJ, Gladman DD, Papp KA, et al. Prevalence of rheumatologist-diagnosed psoriatic arthritis in patients with psoriasis in European/North American dermatology clinics. J Am Acad Dermatol. 2013;69(5):729-735. doi:10.1016/j.jaad.2013.07.023
4. Villani AP, Rouzaud M, Sevrain M, et al. Prevalence of undiagnosed psoriatic arthritis among psoriasis patients: Systematic review and meta-analysis. J Am Acad Dermatol. 2015;73(2):242-248. doi:10.1016/j.jaad.2015.05.001

Contacts: Sun Pharma

Investors:

Dr. Abhishek Sharma
Tel + 91 22 4324 4324, Ext 2929
Tel Direct + 91 22 43242929
Mobile + 91 98196 86016
E mail abhi.sharma@sunpharma.com

Media:

Gaurav Chugh
Tel +91 22 4324 4324, Ext 5373
Tel Direct +91 22 4324 5373
Mobile +91 98104 71414
E mail gaurav.chugh@sunpharma.com

U.S. Corporate:

James Freeman
Mobile: +1 978 808 6956
E-mail: James.Freeman@sunpharma.com